MULTIMEDIA UNIVERSITY

STUDENT ID NO

MULTIMEDIA UNIVERSITY

FINAL EXAMINATION

TRIMESTER 2, 2022/2023

BFS1024 – STATISTICS FOR FINANCE

(All sections / Groups)

7 JULY 2023
9.00 am – 11.00 am
(2 Hours)

INSTRUCTIONS TO STUDENTS

1. This question paper consists of **7 pages** including statistical formulas and statistical tables with **3 questions** only.

2. Attempt **ALL** questions and write your answers in the Answer Booklet provided.

3. Students are allowed to use non-programmable scientific calculators that are permitted to be used in the examination.

Question 1 (25 Marks)

a) In an exclusive restaurant, 3 out of every 5 customers ask for dessert after their meal. A random sample of 10 customers was selected. Find the probability that:
   i. exactly 6 customers ask for dessert after their meal. [4 marks]
   ii. less than 9 customers ask for dessert after their meal. [5 marks]

b) During lunch hours, the arrival of customers at a shopping mall food court is random with an average rate of 90 customers per hour.
   i. What is the probability that exactly two customers will arrive at the food court within a specific one-minute period during lunch hours? [4 marks]
   ii. What is the probability that at least two customers will arrive within a specific two-minute period during lunch hours? [5 marks]

c) The monthly sales of a food and beverage company are approximately normally distributed with a mean of RM 2.5 million and a standard deviation of RM 300,000.
   i. Find the probability that the total sales for a particular month are less than RM 3 million. [3 marks]
   ii. Determine the monthly sales level that has a 5% chance of being exceeded. [4 marks]

Question 2 (30 Marks)

a) The number of calories contained in seven cans of sardines are 190, 189, 188, 192, 187, 190, 191. Assume that the contents are normally distributed.

   i. Calculate the sample mean and standard deviation of the distribution. Leave your answer in 2 decimal places. [5 marks]
   ii. Compute and interpret the 90% and 99% confidence interval estimate of the mean content of calories. [10 marks]
   iii. Based on your answer in (ii), calculate the width of each confidence interval. Describe the effect on the confidence interval estimate of increasing the confidence level. [3 marks]

b) A financial officer for a company wants to estimate the percentage of accounts receivable that are more than 30 days overdue. He surveys 600 accounts and finds that 200 are more than 30 days overdue.
   i. Compute a 90% confidence interval for the true percent of accounts receivable that are more than 30 days overdue and interpret the confidence interval. [7 marks]
   ii. How large is a sample required if the error of the population proportion is within 5% at 95% confidence level? [5 marks]

Continued...

Question 3 (45 Marks)

a) The period of effectiveness of a mosquito repellant produced by a company is approximately normally distributed with a mean of 300 hours and a standard deviation of 30 hours. A sample of 40 repellents was taken and the average period of effectiveness is computed as 290 hours. At 5% significance level, is there evidence that the mean period of effectiveness has changed? Use both test statistics and p-value approach. [13 marks]

b) In the previous year, where a poll was conducted, respondents were asked if they thought unemployment would increase, and 47% thought that it would increase. This year, the poll was conducted again. The result of the poll shows that 284 out of 631 said that they thought unemployment would increase. At the 5% level, is there enough evidence to show that the current proportion of respondents who believe unemployment would increase is less than the proportion in the previous year? Use both test statistics and p-value approach in the hypothesis testing. [14 marks]

c) A manager of a car manufacturing company must decide the type of tires for heavy vehicles that gives better fuel economy. Two types of tires were tested. Type 1 and Type 2. The fuel consumptions in kilometers per litre were recorded as follows:

Truck	Type 1	Type 2
1	4.2	4.1
2	4.7	4.9
3	6.6	6.2
4	7.7	6.9
5	6.7	6.8
6	4.5	4.4
7	5.7	5.7
8	6.0	5.8
9	7.4	6.9
10	4.9	4.7
11	6.1	6.0
12	5.2	4.9

Can the manager infer at 1% significance level that Type 1 tires are better than Type 2 tires in term of fuel consumption? [18 marks]

End of Questions.

FORMULA

A. DESCRIPTIVE STATISTICS

Mean = $\frac{\sum X_i}{n}$

Standard Deviation (s) = $\sqrt{\frac{\sum X^2}{n-1} - \frac{(\sum X)^2}{n(n-1)}}$

B. PROBABILITY

Poisson Probability Distribution

If X follows a Poisson Distribution P (λ) where $P(X = x) = \frac{e^{-\lambda}\lambda^x}{x!}$
then the mean = E(X) = λ and variance = VAR(X) = λ

Binomial Probability Distribution

If X follows a Binomial Distribution B(n, p) where $P(X = x) = {}^nC_x p^x q^{n-x}$

then the mean = E(X) = n p and variance = VAR(X) = npq where q = 1 - p

Normal Distribution

If X follows a Normal distribution N(μ , σ) where E(X) = μ and VAR(X) = σ^2

then $z = \frac{X - \mu}{\sigma}$

C. CONFIDENCE INTERVAL ESTIMATION AND SAMPLE SIZE DETERMINATION

(100 - α) % Confidence Interval for Population Mean (σ Known) = $\overline{X} \pm Z_{\alpha/2}\left(\frac{\sigma}{\sqrt{n}}\right)$

(100 - α)% Confidence Interval for Population Mean (σ Unknown) = $\overline{X} \pm t_{\alpha/2,n-1}\left(\frac{s}{\sqrt{n}}\right)$

(100 - α)% Confidence Interval for Population Proportion = $p \pm Z_{\alpha/2}\sqrt{\frac{p(1-p)}{n}}$

Sample Size Determination for Population Mean $= n \geq \frac{(Z_{\alpha/2})^2\sigma^2}{E^2}$

Continued...

Sample Size Determination for Population Proportion $= n \geq \frac{(Z_{\alpha/2})^2 p(1-p)}{E^2}$

Where E = Limit of Error in Estimation

D. HYPOTHESIS TESTING

One Sample Mean Test

Standard Deviation (σ) Known	**Standard Deviation (σ) Not Known**
$Z = \frac{\bar{x} - \mu}{\sigma/\sqrt{n}}$	$t = \frac{\bar{x} - \mu}{s/\sqrt{n}}$

One Sample Proportion Test

$$Z = \frac{p - \pi}{\sqrt{\frac{\pi(1-\pi)}{n}}}$$

Two Sample Mean Test

Standard Deviation (σ) Known

$$z = \frac{(\overline{x_1} - \overline{x_2})}{\sqrt{\sigma_1^2/n_1 + \sigma_2^2/n_2}}$$

Standard Deviation (σ) Not Known

$$t = \frac{(\overline{x_1} - \overline{x_2})}{\sqrt{S_p^2\left(1/n_1 + 1/n_2\right)}} \quad \textbf{where} \quad S_p^2 = \frac{(n_1 - 1)S_1^2 + (n_2 - 1)S_2^2}{(n_1 - 1) + (n_2 - 1)}$$

Two Sample Proportion Test

$$z = \frac{(p_1 - p_2)}{\sqrt{p(1-p)\left[\frac{1}{n_1} + \frac{1}{n_2}\right]}} \quad \text{where} \quad p = \frac{(n_1 p_1) + (n_2 p_2)}{n_1 + n_2} = \frac{X_1 + X_2}{n_1 + n_2}$$

where X_1 and X_2 are the number of successes from each population

Continued...

TABLE 4. THE NORMAL DISTRIBUTION FUNCTION

The function tabulated is $\Phi(x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{x} e^{-\frac{1}{2}t^2}\, dt$. $\Phi(x)$ is the probability that a random variable, normally distributed with zero mean and unit variance, will be less than or equal to x. When $x < 0$ use $\Phi(x) = 1 - \Phi(-x)$, as the normal distribution with zero mean and unit variance is symmetric about zero.



x	$\Phi(x)$	x	$\Phi(x)$	x	$\Phi(x)$	x	$\Phi(x)$	x	$\Phi(x)$	x	$\Phi(x)$
0·00	0·5000	0·40	0·6554	0·80	0·7881	1·20	0·8849	1·60	0·9452	2·00	0·97725
·01	·5040	·41	·6591	·81	·7910	·21	·8869	·61	·9463	·01	·97778
·02	·5080	·42	·6628	·82	·7939	·22	·8888	·62	·9474	·02	·97831
·03	·5120	·43	·6664	·83	·7967	·23	·8907	·63	·9484	·03	·97882
·04	·5160	·44	·6700	·84	·7995	·24	·8925	·64	·9495	·04	·97932
0·05	0·5199	0·45	0·6736	0·85	0·8023	1·25	0·8944	1·65	0·9505	2·05	0·97982
·06	·5239	·46	·6772	·86	·8051	·26	·8962	·66	·9515	·06	·98030
·07	·5279	·47	·6808	·87	·8078	·27	·8980	·67	·9525	·07	·98077
·08	·5319	·48	·6844	·88	·8106	·28	·8997	·68	·9535	·08	·98124
·09	·5359	·49	·6879	·89	·8133	·29	·9015	·69	·9545	·09	·98169
0·10	0·5398	0·50	0·6915	0·90	0·8159	1·30	0·9032	1·70	0·9554	2·10	0·98214
·11	·5438	·51	·6950	·91	·8186	·31	·9049	·71	·9564	·11	·98257
·12	·5478	·52	·6985	·92	·8212	·32	·9066	·72	·9573	·12	·98300
·13	·5517	·53	·7019	·93	·8238	·33	·9082	·73	·9582	·13	·98341
·14	·5557	·54	·7054	·94	·8264	·34	·9099	·74	·9591	·14	·98382
0·15	0·5596	0·55	0·7088	0·95	0·8289	1·35	0·9115	1·75	0·9599	2·15	0·98422
·16	·5636	·56	·7123	·96	·8315	·36	·9131	·76	·9608	·16	·98461
·17	·5675	·57	·7157	·97	·8340	·37	·9147	·77	·9616	·17	·98500
·18	·5714	·58	·7190	·98	·8365	·38	·9162	·78	·9625	·18	·98537
·19	·5753	·59	·7224	·99	·8389	·39	·9177	·79	·9633	·19	·98574
0·20	0·5793	0·60	0·7257	1·00	0·8413	1·40	0·9192	1·80	0·9641	2·20	0·98610
·21	·5832	·61	·7291	·01	·8438	·41	·9207	·81	·9649	·21	·98645
·22	·5871	·62	·7324	·02	·8461	·42	·9222	·82	·9656	·22	·98679
·23	·5910	·63	·7357	·03	·8485	·43	·9236	·83	·9664	·23	·98713
·24	·5948	·64	·7389	·04	·8508	·44	·9251	·84	·9671	·24	·98745
0·25	0·5987	0·65	0·7422	1·05	0·8531	1·45	0·9265	1·85	0·9678	2·25	0·98778
·26	·6026	·66	·7454	·06	·8554	·46	·9279	·86	·9686	·26	·98809
·27	·6064	·67	·7486	·07	·8577	·47	·9292	·87	·9693	·27	·98840
·28	·6103	·68	·7517	·08	·8599	·48	·9306	·88	·9699	·28	·98870
·29	·6141	·69	·7549	·09	·8621	·49	·9319	·89	·9706	·29	·98899
0·30	0·6179	0·70	0·7580	1·10	0·8643	1·50	0·9332	1·90	0·9713	2·30	0·98928
·31	·6217	·71	·7611	·11	·8665	·51	·9345	·91	·9719	·31	·98956
·32	·6255	·72	·7642	·12	·8686	·52	·9357	·92	·9726	·32	·98983
·33	·6293	·73	·7673	·13	·8708	·53	·9370	·93	·9732	·33	·99010
·34	·6331	·74	·7704	·14	·8729	·54	·9382	·94	·9738	·34	·99036
0·35	0·6368	0·75	0·7734	1·15	0·8749	1·55	0·9394	1·95	0·9744	2·35	0·99061
·36	·6406	·76	·7764	·16	·8770	·56	·9406	·96	·9750	·36	·99086
·37	·6443	·77	·7794	·17	·8790	·57	·9418	·97	·9756	·37	·99111
·38	·6480	·78	·7823	·18	·8810	·58	·9429	·98	·9761	·38	·99134
·39	·6517	·79	·7852	·19	·8830	·59	·9441	·99	·9767	·39	·99158
0·40	0·6554	0·80	0·7881	1·20	0·8849	1·60	0·9452	2·00	0·9772	2·40	0·99180

Continued...

TABLE 4. THE NORMAL DISTRIBUTION FUNCTION

x	$\Phi(x)$	x	$\Phi(x)$	x	$\Phi(x)$	x	$\Phi(x)$	x	$\Phi(x)$	x	$\Phi(x)$
2·40	0·99180	2·55	0·99461	2·70	0·99653	2·85	0·99781	3·00	0·99865	3·15	0·99918
·41	·99202	·56	·99477	·71	·99664	·86	·99788	·01	·99869	·16	·99921
·42	·99224	·57	·99492	·72	·99674	·87	·99795	·02	·99874	·17	·99924
·43	·99245	·58	·99506	·73	·99683	·88	·99801	·03	·99878	·18	·99926
·44	·99266	·59	·99520	·74	·99693	·89	·99807	·04	·99882	·19	·99929
2·45	0·99286	2·60	0·99534	2·75	0·99702	2·90	0·99813	3·05	0·99886	3·20	0·99931
·46	·99305	·61	·99547	·76	·99711	·91	·99819	·06	·99889	·21	·99934
·47	·99324	·62	·99560	·77	·99720	·92	·99825	·07	·99893	·22	·99936
·48	·99343	·63	·99573	·78	·99728	·93	·99831	·08	·99896	·23	·99938
·49	·99361	·64	·99585	·79	·99736	·94	·99836	·09	·99900	·24	·99940
2·50	0·99379	2·65	0·99598	2·80	0·99744	2·95	0·99841	3·10	0·99903	3·25	0·99942
·51	·99396	·66	·99609	·81	·99752	·96	·99846	·11	·99906	·26	·99944
·52	·99413	·67	·99621	·82	·99760	·97	·99851	·12	·99910	·27	·99946
·53	·99430	·68	·99632	·83	·99767	·98	·99856	·13	·99913	·28	·99948
·54	·99446	·69	·99643	·84	·99774	·99	·99861	·14	·99916	·29	·99950
2·55	0·99461	2·70	0·99653	2·85	0·99781	3·00	0·99865	3·15	0·99918	3·30	0·99952

The critical table below gives on the left the range of values of x for which $\Phi(x)$ takes the value on the right, correct to the last figure given; in critical cases, take the upper of the two values of $\Phi(x)$ indicated.

3·075	0·9990	3·263	0·9994	3·731	0·99990	3·916	0·99995
3·105	0·9991	3·320	0·9995	3·759	0·99991	3·976	0·99996
3·138	0·9992	3·389	0·9996	3·791	0·99992	4·055	0·99997
3·174	0·9993	3·480	0·9997	3·826	0·99993	4·173	0·99998
3·215	0·9994	3·615	0·9998	3·867	0·99994	4·417	0·99999
			0·9999		0·99995		1·00000

When $x > 3{\cdot}3$ the formula $1-\Phi(x) \doteq \frac{e^{-\frac{1}{2}x^2}}{x\sqrt{2\pi}}\left[1-\frac{1}{x^2}+\frac{3}{x^4}-\frac{15}{x^6}+\frac{105}{x^8}\right]$ is very accurate, with relative error less than $945/x^{10}$.

TABLE 5. PERCENTAGE POINTS OF THE NORMAL DISTRIBUTION

This table gives percentage points $x(P)$ defined by the equation

$$\frac{P}{100} = \frac{1}{\sqrt{2\pi}}\int_{x(P)}^{\infty} e^{-\frac{1}{2}t^2}\,dt.$$

If X is a variable, normally distributed with zero mean and unit variance, $P/100$ is the probability that $X \geqslant x(P)$. The lower P per cent points are given by symmetry as $-x(P)$, and the probability that $|X| \geqslant x(P)$ is $2P/100$.



P	$x(P)$	P	$x(P)$	P	$x(P)$	P	$x(P)$	P	$x(P)$	P	$x(P)$
50	0·0000	5·0	1·6449	3·0	1·8808	2·0	2·0537	1·0	2·3263	0·10	3·0902
45	0·1257	4·8	1·6646	2·9	1·8957	1·9	2·0749	0·9	2·3656	0·09	3·1214
40	0·2533	4·6	1·6849	2·8	1·9110	1·8	2·0969	0·8	2·4089	0·08	3·1559
35	0·3853	4·4	1·7060	2·7	1·9268	1·7	2·1201	0·7	2·4573	0·07	3·1947
30	0·5244	4·2	1·7279	2·6	1·9431	1·6	2·1444	0·6	2·5121	0·06	3·2389
25	0·6745	4·0	1·7507	2·5	1·9600	1·5	2·1701	0·5	2·5758	0·05	3·2905
20	0·8416	3·8	1·7744	2·4	1·9774	1·4	2·1973	0·4	2·6521	0·01	3·7190
15	1·0364	3·6	1·7991	2·3	1·9954	1·3	2·2262	0·3	2·7478	0·005	3·8906
10	1·2816	3·4	1·8250	2·2	2·0141	1·2	2·2571	0·2	2·8782	0·001	4·2649
5	1·6449	3·2	1·8522	2·1	2·0335	1·1	2·2904	0·1	3·0902	0·0005	4·4172

Continued...

TABLE 10. PERCENTAGE POINTS OF THE t-DISTRIBUTION

This table gives percentage points $t_\nu(P)$ defined by the equation

$$\frac{P}{100} = \frac{1}{\sqrt{\nu\pi}} \frac{\Gamma(\frac{1}{2}\nu+\frac{1}{2})}{\Gamma(\frac{1}{2}\nu)} \int_{t_\nu(P)}^{\infty} \frac{dt}{(1+t^2/\nu)^{\frac{1}{2}(\nu+1)}}.$$

Let X_1 and X_2 be independent random variables having a normal distribution with zero mean and unit variance and a χ^2-distribution with ν degrees of freedom respectively; then $t = X_1/\sqrt{X_2/\nu}$ has Student's t-distribution with ν degrees of freedom, and the probability that $t \geqslant t_\nu(P)$ is $P/100$. The lower percentage points are given by symmetry as $-t_\nu(P)$, and the probability that $|t| \geqslant t_\nu(P)$ is $2P/100$.



The limiting distribution of t as ν tends to infinity is the normal distribution with zero mean and unit variance. When ν is large interpolation in ν should be harmonic.

P	40	30	25	20	15	10	8	2·5	2	0·5	0·2	0·05
$\nu = 1$	0·3249	0·7265	1·0000	1·3764	1·963	3·078	6·314	12·71	31·82	63·66	318·3	636·6
2	0·2887	0·6172	0·8165	1·0607	1·386	1·886	2·920	4·303	6·965	9·925	22·33	31·60
3	0·2767	0·5844	0·7649	0·9785	1·250	1·638	2·353	3·182	4·541	5·841	10·21	12·92
4	0·2707	0·5686	0·7407	0·9410	1·190	1·533	2·132	2·776	3·747	4·604	7·173	8·610
5	0·2672	0·5594	0·7267	0·9195	1·156	1·476	2·015	2·571	3·365	4·032	5·893	6·869
6	0·2648	0·5534	0·7176	0·9057	1·134	1·440	1·943	2·447	3·143	3·707	5·208	5·959
7	0·2632	0·5491	0·7111	0·8960	1·119	1·415	1·895	2·365	2·998	3·499	4·785	5·408
8	0·2619	0·5459	0·7064	0·8889	1·108	1·397	1·860	2·306	2·896	3·355	4·501	5·041
9	0·2610	0·5435	0·7027	0·8834	1·100	1·383	1·833	2·262	2·821	3·250	4·297	4·781
10	0·2602	0·5415	0·6998	0·8791	1·093	1·372	1·812	2·228	2·764	3·169	4·144	4·587
11	0·2596	0·5399	0·6974	0·8755	1·088	1·363	1·796	2·201	2·718	3·106	4·025	4·437
12	0·2590	0·5386	0·6955	0·8726	1·083	1·356	1·782	2·179	2·681	3·055	3·930	4·318
13	0·2586	0·5375	0·6938	0·8702	1·079	1·350	1·771	2·160	2·650	3·012	3·852	4·221
14	0·2582	0·5366	0·6924	0·8681	1·076	1·345	1·761	2·145	2·624	2·977	3·787	4·140
15	0·2579	0·5357	0·6912	0·8662	1·074	1·341	1·753	2·131	2·602	2·947	3·733	4·073
16	0·2576	0·5350	0·6901	0·8647	1·071	1·337	1·746	2·120	2·583	2·921	3·686	4·015
17	0·2573	0·5344	0·6892	0·8633	1·069	1·333	1·740	2·110	2·567	2·898	3·646	3·965
18	0·2571	0·5338	0·6884	0·8620	1·067	1·330	1·734	2·101	2·552	2·878	3·611	3·922
19	0·2569	0·5333	0·6876	0·8610	1·066	1·328	1·729	2·093	2·539	2·861	3·579	3·883
20	0·2567	0·5329	0·6870	0·8600	1·064	1·325	1·725	2·086	2·528	2·845	3·552	3·850
21	0·2566	0·5325	0·6864	0·8591	1·063	1·323	1·721	2·080	2·518	2·831	3·527	3·819
22	0·2564	0·5321	0·6858	0·8583	1·061	1·321	1·717	2·074	2·508	2·819	3·505	3·792
23	0·2563	0·5317	0·6853	0·8575	1·060	1·319	1·714	2·069	2·500	2·807	3·485	3·768
24	0·2562	0·5314	0·6848	0·8569	1·059	1·318	1·711	2·064	2·492	2·797	3·467	3·745
25	0·2561	0·5312	0·6844	0·8562	1·058	1·316	1·708	2·060	2·485	2·787	3·450	3·725
26	0·2560	0·5309	0·6840	0·8557	1·058	1·315	1·706	2·056	2·479	2·779	3·435	3·707
27	0·2559	0·5306	0·6837	0·8551	1·057	1·314	1·703	2·052	2·473	2·771	3·421	3·690
28	0·2558	0·5304	0·6834	0·8546	1·056	1·313	1·701	2·048	2·467	2·763	3·408	3·674
29	0·2557	0·5302	0·6830	0·8542	1·055	1·311	1·699	2·045	2·462	2·756	3·396	3·659
30	0·2556	0·5300	0·6828	0·8538	1·055	1·310	1·697	2·042	2·457	2·750	3·385	3·646
32	0·2555	0·5297	0·6822	0·8530	1·054	1·309	1·694	2·037	2·449	2·738	3·365	3·622
34	0·2553	0·5294	0·6818	0·8523	1·052	1·307	1·691	2·032	2·441	2·728	3·348	3·601
36	0·2552	0·5291	0·6814	0·8517	1·052	1·306	1·688	2·028	2·434	2·719	3·333	3·582
38	0·2551	0·5288	0·6810	0·8512	1·051	1·304	1·686	2·024	2·429	2·712	3·319	3·566
40	0·2550	0·5286	0·6807	0·8507	1·050	1·303	1·684	2·021	2·423	2·704	3·307	3·551
50	0·2547	0·5278	0·6794	0·8489	1·047	1·299	1·676	2·009	2·403	2·678	3·261	3·496
60	0·2545	0·5272	0·6786	0·8477	1·045	1·296	1·671	2·000	2·390	2·660	3·232	3·460
120	0·2539	0·5258	0·6765	0·8446	1·041	1·289	1·658	1·980	2·358	2·617	3·160	3·373
∞	0·2533	0·5244	0·6745	0·8416	1·036	1·282	1·645	1·960	2·326	2·576	3·090	3·291

End of Pages.